

# Advantages and Disadvantages of Aquatic Plant Control Techniques

John D. Madsen

US Dept. Agriculture – Agricultural Research Service  
Exotic and Invasive Weed Research Unit, Davis, CA  
[jmadsen@ucdavis.edu](mailto:jmadsen@ucdavis.edu)

2014 MI Inland Lakes Convention, May 1, 2014, Boyne Falls, MI

# Aquatic Plant Management Plan

- Prevention
- Assessment
- Site-specific management
- Evaluation
- Monitoring
- Education



Hydrilla topped out in Lake Guntersville, AL – in 12' water depth.  
Photo by RM Stewart.

# Prevention



- Prevent invasive species introductions
  - Signage
  - Regulations
  - Enforcement
  - Monitor lakes
- Prevent eutrophication and sedimentation

# Environmental Impacts of Aquatic Plant Management

- All have environmental impacts on nontarget species and ecosystem processes
- Impacts are both direct and indirect
- Impacts have not been determined for some techniques
- Public perceptions not supported by existing knowledge

# Unequal Perceptions of Techniques

- “Chemicals are ‘poisons’ put into the water; therefore are bad”
- “Harvesting is just like cutting your lawn, so it isn’t bad”
- “Dredging projects ruin natural substrates and are bad”
- “Biological control agents are natural, so they are good”

# Direct Effects

- Specific and direct mechanism of effect
  - Grass carp eats plants (target or nontarget)
  - Herbicide kills plants (target or nontarget)
  - Harvester harvests fish, macroinvertebrates and plants

# Indirect Effects

- Operates through a separate step or process
  - Dredging deepens the water, making it too deep for plants
  - Herbicides kill plants, which reduces refugia for fish
  - Harvesters stir up sediment, which causes an algal bloom

# Aquatic Plant Control Techniques

- Biological Control
- Chemical Control
- Mechanical Control
- Physical Control
- “Institutional Control”
- “No (Active) Control”





# Aquatic Plant Management References

Gettys, LA, WT Haller and M Bellaud, eds. 2009. Biology and control of aquatic plants: a best management practices handbook. Aquatic Ecosystem Restoration Foundation, Marietta GA. 210 pages. [www.aquatics.org](http://www.aquatics.org)

Madsen, J. D. 1997. Ch. 12. Method for Management of Nonindigenous Aquatic Plants, pp. 145-171. In: J. O. Luken and J. W. Thieret, eds. Assessment and Management of Plant Invasions. Springer, New York. 316pp. ISBN 0-387-94809-0

Madsen, J.D. 2000. Advantages and Disadvantages of Aquatic Plant Management Techniques. ***US Army Engineer Research and Development Center Miscellaneous Report ERDC/EL MP-00-1***, Vicksburg, MS. September 2000.

# Potential Environmental Impacts

- Water quality
- Plankton
- Nontarget plants
- Macroinvertebrates
- Fish
- Waterfowl
- Human health, welfare, and utilization



# Biological Control



## Insects

- Classical
- Native

## Grass Carp

## Pathogens

- Classical
- Native

## Plant Restoration

# Grass Carp

- Advantages
  - Effective
  - Inexpensive
  - Long-term
- Disadvantages
  - “All-or-none” response
  - Not selective
  - Cannot control feeding sites
  - Cannot stop fish
  - Difficult to contain
  - Reproduction?
  - Won't eat some species (e.g., Eurasian watermilfoil)
  - Regulations



# Environmental Effects of Grass Carp

Abstract.—We investigated effects of triploid grass carp *Ctenopharyngodon idella* on aquatic macrophyte communities, water quality, and public satisfaction for 98 lakes and ponds in Washington State stocked with grass carp between 1990 and 1995. Grass carp had few noticeable effects on macrophyte communities until 19 months following stocking. After 19 months, submersed macrophytes were either completely eradicated (39% of the lakes) or not controlled (42% of the lakes) in most lakes. Intermediate control of submersed macrophytes occurred in 18% of lakes at a median stocking rate of 24 fish per vegetated surface acre. Most of the landowners interviewed (83%) were satisfied with the results of introducing grass carp. For sites where all submersed macrophytes were eradicated, average turbidity was higher (11 nephelometric turbidity units, NTU) than at sites where macrophytes were controlled to intermediate levels (4 NTU) or unaffected by grass carp grazing (5 NTU).

Bonar, S.A., B. Boldin, and M. Divens. 2002. Effects of triploid grass carp on aquatic plants, water quality, and public satisfaction in Washington State. *N. Am. J. Fish. Manage.* 22:96-105.

# Classical Insect Control

- Advantages
  - Public perception
  - Low cost after R&D
  - Long-term
  - Works well for some species in some areas
- Disadvantages
  - No agents for several target invasive plants
  - Long time for R&D
  - Unpredictability of results
  - Limited distribution of effectiveness



*Agasicles hygrophila*  
Alligatorweed flea beetle  
Copyright 1997 USDA-ARS

# Native Insect Biocontrol

- Advantages
  - Public perception
  - Avoid quarantine period and problems
  - Several apparent successes through natural populations
- Disadvantages
  - Not an operational technique at this time
  - Unpredictability of results
  - No current strategy for use
  - Very expensive
  - No theoretical basis for long-term success



# Fungal Pathogens

- Advantages

- Typical plant diseases that can be effective
- Can be combined with other techniques

- Disadvantages

- No operational formulations
- To date, only a “contact mycoherbicide” is under R&D



# Native Plant Restoration

- Advantages
  - Provides habitat for aquatic organisms
  - May retard reinvasion
  - Public perception
- Disadvantages
  - Not a control technique
  - Restorative
  - Techniques and approach largely under R&D
  - Very labor intensive



# Chemical Control



## Contact

- Carfentrazone ethyl
- Complexed copper
- Diquat
- Endothall
- Flumioxazin


## Systemic

- 2,4-D
- Bispyribac-sodium
- Fluridone
- Glyphosate\*
- Imazamox
- Imazapyr\*
- Penoxsulam
- Triclopyr


Some would also include sodium carbonate peroxyhydrate (an algaecide) and water dyes (alters water transparency)

# Aquatic Applications of Herbicides

- Herbicides are applied to water, and plants take up from water
- Water movement, residence time, and concentration critical to effectiveness



# Concentration / Exposure Time



# Contact Herbicides

- Advantages

- Requires only a short exposure time (6-24 hours)
- Broad spectrum
- Very fast action (Usually less than 7 days)
- Inexpensive (~\$200/acre)

- Disadvantages

- Does not kill entire plant
- Requires more than one treatment per year
- Not selective
- Public perception

# Contact Herbicide Impact

A nearly monotypic population of egeria (*Egeria densa* Planch.) was interfering with recreation in a popular lake in southwest Washington. In June 2003 the littoral zone was treated with the contact herbicide diquat (6,7-dihydrodipyrido [1,2-a2',1'-c] pyrazinediium dibromide). Aquatic plant frequency and biomass data were collected on all submersed species before treatment, and eight weeks, one year and two years after treatment. Water quality and herbicide dissipation data were also collected before and for one season after the treatment. Results from the aquatic plant data showed a significant reduction in egeria frequency and biomass after the herbicide treatment, although the species did not disappear entirely. **Two native submersed species, water moss (*Fontinalis antipyretica* Hedw.) and stonewort (*Nitella* sp.), increased after treatment.** However their increase was not enough to offset the egeria population reduction, as total plant abundance was significantly reduced after treatment. The herbicide dissipation data illustrated the dispersal of diquat throughout the lake and persistence at low concentrations (up to 10.4 ppb) in the water column for at least two weeks after treatment. **Water quality data demonstrated a slight decrease in dissolved oxygen and water transparency following the herbicide treatment, potentially due to plant die-off and subsequent plant decomposition.**

Parsons, J.K., K.S. Hamel, and R. Wierenga. 2007. J. Aquat. Plant Manage. 45:35.

# Systemic - 2,4-D

- Advantages
  - Selective
    - Eurasian watermilfoil
  - Intermediate exposure time
  - Inexpensive (~\$300/acre)
  - Usually effective
  - Moderate response time (1-2 weeks)
- Disadvantages
  - Public perception
  - Does not work on some plants
  - Effective for 1 to 3 years



# Couch and Nelson. 1982. JAPM 20:8- 13

The butoxyethanol ester formulation of the herbicide 2,4-dichlorophenoxyacetic acid (2,4-D BEE) was applied for the control of Eurasian watermilfoil (*Myriophyllum spicatum* L.) in Kerr Reservoir in 1977 and 1978. The only detrimental effect to non-target species that could be attributed to herbicide treatment was a short-term depression of phytoplankton populations in those sites sampled in 1977 where large areas of the reservoir were treated or when the treated areas were in places sheltered from the action of currents and waves. Therefore, within the restraints and conditions of this investigation, it may be concluded that 2,4-D did not adversely affect the non-target components in the sampled ecosystems.



# Whitney et al. 1973.

## JAPM 11:13-17.

On June 4, 1968 an application of a 20% (acid equivalent) formulation of the butoxyethanol ester of (2,4-dichlorophenoxy)acetic acid (2,4-D) herbicide was conducted to 200 acres of Eurasian watermilfoil (*Myriophyllum spicatum* L.) in Currituck Sound, North Carolina. No acute adverse effects on fish and other organisms were observed. No water samples contained residues exceeding 0.10 ppm. Reduction of Eurasian watermilfoil was estimated at 95% and subsequent re-establishment of native plants was considered to be of significant benefit to waterfowl. It is conceivable that reinfestation by Eurasian watermilfoil could have been reduced and possibly prevented through total treatment of the Sound.

# Systemic - Fluridone

- Advantages
  - Broad spectrum
  - Very low concentration (5-25 ppm)
  - Moderately expensive (~\$600/acre)
  - No drinking or contact restrictions
- Disadvantages
  - Selective only at very low rates
  - Very long exposure time (30-60 days)
  - Very long plant response time (30-90 days)




# Mechanical Control



- Hand pulling
- Cutting
- Harvesting
- Diver-operated suction harvesting
- Rotovating

# Hand Harvesting

- Advantages
  - Low technology
  - Widely used in many parts of the world
  - Effective in small areas
  - Can be selective
  - Affordable (volunteers)
- Disadvantages
  - Very labor intensive
  - Not practical for large areas (<<1 acre)



# Cutting / Harvesting

- Advantages
  - Direct relief
  - Immediate efficacy
  - Moderately expensive (~\$400/acre)
  - Public perception
- Disadvantages
  - Not selective
  - Short-term control (2-3 harvests per season)
  - May aid spread some species
  - Slow
  - Disposal (?)

# Total Fish Impact: Largemouth Bass

Mikol, G.F. 1985. Journal of Aquatic Plant Management 23:59-63.

TABLE 4. SUMMARY OF DIRET EFFECTS OF 1982 MECHANICAL HARVESTING ON JUVENILE LARGEMOUTH BASS (*Micropterus salmoides*).

	COLLECTION DATE		
	8/13/82 <sup>1</sup>		
	Site #1	Site #2	Combined
Total # fish removed	11	7	18
# fish removed/ha	220	56	103
Fish standing crop estimate (#/ha)	1,894	1,894	1,894
% standing crop removed	11.6	3.0	5.4

<sup>1</sup>Site #1 was previously harvested in June 1982. Site #2 was previously unharvested.

# Diver-operated Suction Harvester

- Advantages
  - Selective (dependent on operator and environment)
  - Longer-term control
- Disadvantages
  - Very limited areas
  - Very slow
  - Expensive (~\$1,000/acre?)
  - Disposal (?)

# Rotovating

- Advantages
  - Longer term than other mechanical (on Eurasian watermilfoil)
  - Moderately inexpensive
- Disadvantages
  - Turbidity
  - Spreads fragments
  - Limited environmental range by depth, sediment
  - Free-floating plant material





# Physical Control



- Dredging
- Drawdown
- Benthic Barrier
- Shading
- Nutrient Inactivation

# Dredging

- Advantages

- Very effective
- Very long term

- Disadvantages

- Very expensive (~\$6,000/acre)
- Not selective
- Impacts on other organisms?
- Dredge spoils

# Drawdown

- Advantages

- Effective
- Very inexpensive (~\$0/acre?)
- Moderate-term

- Disadvantages

- Not selective
- Impacts on other organisms (?)
- Impacts on human uses
- Need water control structure

# Drawdown and Nutrient Loading

Studies on lake sediments showed that drying and freezing, as would occur during the management practice of water level drawdown, increased the release of phosphorus from the sediments over controls. ***P release from sediments increased as a result of drying in all ten lake sites studied***, but the amount of P released varied significantly between sites. ... Freezing of sediments also resulted in increased nutrient release, with 70 times more P released from frozen sediments compared to unfrozen controls. The full effect of freezing was realized within 3 days. The combined effect of drying then freezing did not increase the amount of P released above that of sediments that were frozen without prior drying. P release as a result of drying was significantly correlated with sediment P and organic content. The data indicate that the P released resulted from the death of microorganisms due to drying and freezing. ***The results show that lake drawdown may significantly increase internal P loading to lakes of central New York State.***

Klotz, R.L. and S.A. Linn. 2001. Lake Reserv. Manage. 17:48-54.

# Benthic Barrier

- Advantages

- Effective
- Broad spectrum
- Immediate effect
- Moderate term (several years)

- Disadvantages

- Small scale
- Expensive (~\$3,000/acre)
- Impacts on other organisms, fish spawning
- Not selective

# Institutional Control

- Quarantine
- Regulation
- Prevention
- Watershed management



# Institutional Control

- Advantages

- May prevent problems from happening
- May improve water quality over the long term
- Relatively inexpensive (depending on solution)

- Disadvantages

- May not redress current problems
- Does little to alleviate short-term problems
- May require cooperation and involvement of many diverse groups

# “No Control”

Shows a lot of hope:

- Hope the plants go away
- Hope the people asking for permits go away
- Hope the people asking for money go away





## In other words...

- The problem (invasive aquatic plants like hydrilla and Eurasian watermilfoil) are worse than the potential solutions; that's why we would manage these plants.
- While the solutions are not without some environmental downside, the drawbacks are less than doing nothing

# Environmental Impact

- All techniques have positive and negative environmental attributes
- Evaluate true environmental impacts of all techniques, not “public perceptions”
- Emphasize quantitative evaluation of all costs and benefits
- It is time for a balanced assessment of ecosystem risks


# Acknowledgements



I thank the Aquatic Ecosystem Restoration Foundation, who kindly provided this opportunity for me to visit Michigan in the springtime

# Contact Information

Dr. John D. Madsen  
USDA-ARS, EIWRU  
Univ. California-Davis  
Dept. of Plant Sciences, MS 4  
One Shields Ave.  
Davis, CA 95616  
Ph. 530-752-7870  
Fax 530-752-4604  
E-mail: [jmadsen@ucdavis.edu](mailto:jmadsen@ucdavis.edu)



John Madsen applying herbicides to bareground plots with flowering rush, in Pend Oreille Lake, ID. Photo by Tom Woolf